

ZETA FUNCTIONS OF INFINITE GRAPH BUNDLES

SAMUEL COOPER* AND STRATOS PRASSIDIS**

ABSTRACT. We compute the equivariant zeta function for bundles over infinite graphs and for infinite covers. In particular, we give a “transfer formula” for the zeta function of infinite graph covers. Also, when the infinite cover is given as a limit of finite covers, we give a formula for the limit of the zeta functions.

1. INTRODUCTION

The Ihara zeta function of a finite graph reflects combinatorial and spectral properties of that graph ([13], [2], [18]). Originally, Ihara defined the zeta function on finite graphs imitating the classical definition of the zeta function:

$$\zeta_X(z) = \prod_{[C]} (1 - z^{\ell(C)})^{-1},$$

where the product is over all equivalence classes of primitive closed loops C in X and $\ell(C)$ denotes the length of C . In [2], it was shown that, for a finite graph X :

$$\zeta_X(z)^{-1} = (1 - z^2)^{\epsilon - \nu} \det(I - zA + z^2Q),$$

where ϵ is the number of edges, ν is the number of vertices, A is the adjacency matrix of X , and Q is the diagonal matrix with entries $\deg(v) - 1$, for each $v \in V(X)$. In [8], the definition of Ihara zeta function was extended to infinite graphs that are limits of sequences of finite graphs. In particular, it was shown in [8], using the results in [17], that the sequence of the zeta functions of the finite graphs converges. In [3], [4], [10], [11], [12], the expression of the zeta function as a rational function was extended to infinite graphs that admit an action of a discrete group Γ with finite quotient. The determinant in the finite case is replaced by the determinant in a von Neumann algebra $\mathcal{N}_0(X, \Gamma)$ of all the bounded operators on $L^2(V(X))$. In [6] the zeta function of finite graph bundles over finite was computed generalizing the results on graph coverings that appear [18], [19], [20]. Their results can be described as transfer results for the Ihara zeta function.

We combine the results on infinite graphs and bundles to derive a transfer formula for infinite bundles and coverings. Let ϕ be an $\text{Aut}(F)$ -assignment on X . Let (Γ, Δ) be a pair of groups that act on X and F in such a way that the actions are ϕ -compatible and by finite co-volume.

2000 *Mathematics Subject Classification.* Primary 11M36; Secondary 05C50, 11M41.

*Partially supported by an NSF REU grant.

**Partially supported by Canisius College Summer Grant and an NSF REU grant.

Theorem (Main Theorem 1). *With the above assumptions, the equivariant zeta function is given by:*

$$\zeta_{X \times^\psi F, \Gamma \times \Delta}(z)^{-1} = (1 - z^2)^{-\chi^{(2)}(X \times^\psi F)} \det_{\Gamma \times \Delta} \left(I - \sum_{\gamma \in \text{Aut}(F)} (A_{\vec{X}_{(\psi, \gamma)}} \otimes P_\gamma + I_X \otimes A_F) z + Q z^2 \right),$$

where $A_{\vec{X}_{(\psi, \gamma)}}$ is the adjacency matrix of the directed graph spanned by the edges in $\psi^{-1}(\gamma)$, P_γ is the permutation matrix induced by the action of γ on $V(F)$, $\chi^{(2)}$ is the Euler characteristic of the quotient $X_1 \times^{\phi_1} F$, $\det_{\Gamma \times \Delta}$ is the determinant defined on the von Neumann algebra of $\Gamma \times \Delta$, and Q is the diagonal operator such that $Q(y, i) = \deg(y) + \deg(i) - 1$.

Using similar methods, we prove a decomposition formula of the Ihara zeta function for infinite covers. Let $p : Y \rightarrow X$ be a cover with X finite. Let $\Gamma = \text{Cov}(p)$.

Theorem (Main Theorem 2). *With the above notation,*

$$\zeta_{Y, \Gamma}(z)^{-1} = (1 - z^2)^{-\chi^{(2)}(Y)} \det_\Gamma \left(I - \left(\sum_{\gamma \in \Gamma} A_{(\vec{Y}_{(\psi, \gamma)})} \otimes P_\gamma \right) z + Q z^2 \right),$$

where Q is the diagonal operator with (x, γ) -entry $\deg(x) - 1$.

We apply the above calculations to sequences of *strongly convergent graphs*. In particular, a sequence $\{(X_n, w_n)\}_{n \in \mathbb{N}}$ is strongly convergent to (X, w) if it is a covering sequence of regular graphs converging to X in such a way that X covers compatibly each element of the sequence. Such sequences appear when we consider the Cayley graphs of finite quotients of a groups converging to the Cayley graph of the group.

2. PRELIMINARIES

We now define a number of terms that we will use later on.

Definition 2.1. Let G be any locally finite graph. Then we define the adjacency operator A_G as follows: for any $u, v \in V(G)$,

$$A_G(u, v) = \begin{cases} 1, & \text{if } u \sim v \\ 0, & \text{otherwise.} \end{cases}$$

The definition makes sense even if the graph is directed. If G is undirected, the A_G is symmetric.

Definition 2.2. Let $\tilde{G}$ and G be locally finite graphs. We say that

$$p : \tilde{G} \rightarrow G$$

is a graph covering if the following two conditions hold:

- (1) If $x \sim_{\tilde{G}} y$, then $p(x) \sim_G p(y)$.

- (2) For any $x \in \tilde{G}$, $p : N(x) \rightarrow N(p(x))$ is a bijection.

The first condition in the definition means that p is a graph map. The second condition is a local triviality condition.

Graph bundles are defined in [15]. They generalize the graph coverings in the sense that the “fiber graph” is allowed to have a non-empty set of edges. We will concentrate on bundles with finite fibers. For a graph X , we denote by $E(\vec{X})$ the set of ordered edges—i.e., each edge of X appears twice, each with opposite orientation.

Definition 2.3. Let G be any locally finite graph (possibly infinite), let F be a finite graph. We define an $\text{Aut}(F)$ -voltage assignment on G by

$$\phi : E(\vec{G}) \rightarrow \text{Aut}(F), \quad \phi(uv) = \phi(vu)^{-1}.$$

Definition 2.4. Let G be a locally finite graph, F a finite graph, and ϕ an $\text{Aut}(F)$ -voltage assignment on G . We define a graph bundle $G \times^\phi F$ to be the graph with vertex set $V(G) \times V(F)$, with two vertices $(u, i), (v, j) \in G \times^\phi F$ adjacent if either one of the following two conditions hold:

- (1) $u \sim v$ and $j = i^{\phi(uv)}$
- (2) $u = v$ and $i \sim j$.

Let ϕ be a $\text{Aut}(F)$ -voltage assignment on G . Let $\gamma \in \text{Aut}(F)$.

- (1) Let $\vec{G}_{(\phi, \gamma)}$ denote the spanning subgraph of the digraph $\vec{G}$ whose directed edge set is $\phi^{-1}(\gamma)$.
- (2) We define the permutation operator P_γ by the following formula: for any two vertices i, j in $V(F)$,

$$P_\gamma(i, j) = \begin{cases} 1, & \text{if } j = i^\gamma \\ 0, & \text{otherwise.} \end{cases}$$

Remark 2.5. When the graphs are infinite, the matrices defined above are operators on the Hilbert space with basis the vertex set of the graph. More precisely, if G is any locally finite graph, set $L^2(G)$ to be the Hilbert space:

$$L^2(G) = \left\{ f : V(G) \rightarrow \mathbb{C} : \sum_{u \in V(G)} |f(u)|^2 < \infty \right\}.$$

Then the adjacency operator is given by

$$A(f)(u) = \sum_{u \sim v} f(v).$$

With the above notation,

$$P_\gamma(f)(i) = f(\gamma(i)).$$

The following combines covering maps and bundles.

Theorem 2.6. *Let F and X be locally finite graphs. Let X be equipped with an $\text{Aut}(F)$ -voltage assignment ϕ . Let $p : Y \rightarrow X$ be a covering map and ψ the $\text{Aut}(F)$ -voltage assignment*

$$\psi : E(\vec{Y}) \rightarrow \text{Aut}(F), \quad \psi(xy) = \phi(p(x)p(y)).$$

Define a graph map

$$\tilde{p} : Y \times^\psi F \rightarrow X \times^\phi F, \quad \tilde{p}(x, i) = (p(x), i).$$

Then $\tilde{p}$ is a covering map.

Proof. First we will show prove that $\tilde{p}$ is a graph map i.e., that is preserves adjacency. Let $(x, i) \sim (y, j)$ in $Y \times^\psi F$. There are two cases to consider:

- (1) Suppose $x \sim y$ in Y . Then $p(x) \sim p(y)$, and $j = i^{\psi(xy)} = i^{\phi(p(x)p(y))}$. Thus, by definition,

$$\tilde{p}(x, i) = (p(x), i) \sim (p(y), j) = \tilde{p}(y, j), \quad \text{in } X \times^\phi F.$$

- (2) Suppose $x = y$. Then $i \sim j$ in F , and clearly $p(x) = p(y)$. Thus, by definition, $\tilde{p}(x, i) \sim \tilde{p}(y, j)$.

Thus $\tilde{p}$ preserves adjacency.

Now we must show that $\tilde{p}|_{N(x, i)}$ is a bijection.

$\tilde{p}|_{N(x, i)}$ is an injection. Let $(y_1, j_1), (y_2, j_2) \in N(x, i)$ with $\tilde{p}(y_1, j_1) = \tilde{p}(y_2, j_2)$. Then we know that $p(y_1) = p(y_2)$ and $j_1 = j_2$. Now there are two cases to consider:

- (1) Suppose $y_1 = x$ and $i \sim j_1$. Then $i \sim j_2$, and since $(x, i) \sim (y_2, j_2)$, we must have $x = y_2$. Thus $y_1 = y_2$, so $(y_1, j_1) = (y_2, j_2)$. The same argument works if $y_2 = x$.
- (2) Suppose $y_1 \sim x$, and $i = j_1^{\psi(y_1 x)} = j_1^{\phi(p(y_1)p(x))}$. Since $p(y_1) = p(y_2)$, we see that $i = j_2^{\phi(p(y_2)p(x))}$. Thus, since $(y_2, j_2) \sim (x, i)$, we must have $y_2 \sim x$. Now, since p is a graph covering map, $p|_{N(x)}$ is a bijection. But $y_1, y_2 \in N(x)$ and $p(y_1) = p(y_2)$; thus, $y_2 = y_1$, so $(y_1, j_1) = (y_2, j_2)$.

$\tilde{p}|_{N(x, i)}$ is a surjection. Let $(u, k) \in N((p(x), i))$. Again there are two cases to consider:

- (1) Suppose $u = p(x)$ and $i \sim k$. Then $\tilde{p}(x, i) = (u, k)$, and by definition, $(x, i) \in N(x, i)$.
- (2) Suppose $u \sim p(x)$ and $i = k^{\phi(up(x))}$. Since $p|_{N(x)}$ is a surjection, there exists some $y \in N(x)$ such that $p(y) = u$. Then $y \sim x$ and $i = k^{\phi(p(y)p(x))} = k^{\psi(yx)}$. Thus, $(y, k) \in N(x, i)$ and $\tilde{p}(y, k) = (u, k)$.

Therefore, $\tilde{p}|_{N(x, i)}$ is a bijection. This completes the proof. $\square$

The vertex set of a bundle over G is $V(G) \times V(F)$. Then

$$L^2(G \times^\phi F) = L^2(G) \otimes L^2(F)$$

where the tensor product takes place in the category of Hilbert spaces. More precisely, it is the completion of the algebraic tensor product. The following theorem (proved in [15] for the finite case) provides a decomposition for the adjacency operator of any graph bundle.

Theorem 2.7. *Let ϕ be an $\text{Aut}(F)$ -voltage assignment on a locally finite graph G , with F locally finite. Then*

$$A_{G \times^\phi F} = \bigoplus_{\gamma \in \text{Aut}(F)} A_{\vec{G}_{(\phi, \gamma)}} \otimes P_\gamma + I_G \otimes A_F.$$

Proof. It is enough to prove the result for functions of the form $f \otimes g$, where $f \in L^2(G)$ and $g \in L^2(F)$. Let $(u, i) \in V(G \times^\phi F)$. Then

$$A_{G \times^\phi F}(f \otimes g)(u, i) = \sum_{(u, i) \sim (v, j)} f(v)g(j).$$

The right hand side is given by:

$$\bigoplus_{\gamma \in \text{Aut}(F)} A_{\vec{G}_{(\phi, \gamma)}} \otimes P_\gamma(f \otimes g)(u, j) + I_G \otimes A_F(f \otimes g)(u, j) = \bigoplus_{\gamma \in \text{Aut}(F), u \in \vec{G}_{(\phi, \gamma)}} A_{\vec{G}_{(\phi, \gamma)}}(f|)(u)P_\gamma(g)(i) + f(u)A_F(g)(i)$$

There are two possibilities for $(u, i) \sim (v, j)$:

- (1) $u \sim v$ and $i = j^{\phi(uv)}$. Then the right hand side becomes:

$$A_{\vec{G}_{(\phi, \phi(uv))}}(f|)(u)P_\gamma(g)(i) = f(v)g(j).$$

- (2) $u = v$ and $i \sim j$. In the right hand side, only the last summand is non-zero and it is equal to $f(u)g(j)$.

Finally, it is clear that if neither $u \sim v$ nor $u = v$, then the sum on the right hand side is zero. This completes the proof. $\square$

By a *marked graph*, we mean a pair (X, w) with X a graph and w a distinguished vertex.

Definition 2.8. On the space of marked graphs there is a metric dist defined as follow:

$$\text{dist}\left((X_1, w_1), (X_2, w_2)\right) = \inf \left\{ \frac{1}{n+1}; B_{X_1}(w_1, n) \text{ is isometric to } B_{X_2}(w_2, n) \right\},$$

where $B_X(w, n)$ is the combinatorial ball of radius n in X centered on w .

For a sequence of marked graphs $\{(X_n, w_n)\}_{n \in \mathbb{N}}$, we say that (X, w) is the limit graph if

$$\lim_{n \rightarrow \infty} \text{dist}((X, w), (X_n, w_n)) = 0.$$

For a finite graph X , let $c_r(X)$ denote the number of closed paths in X of length r . Let

$$(X, w) = \lim_{n \rightarrow \infty} (X_n, w_n),$$

where $\{(X_n, w_n)\}_{n \in \mathbb{N}}$ is a covering sequence of k -regular marked graph. In [8], the definition of the number c_r is extended for the graph X as follows:

$$\tilde{c}_r = \lim_{n \rightarrow \infty} \frac{c_r(X_n)}{|X_n|}.$$

In [8], it was shown that the limit exists. The zeta function $\zeta(X, w)$ of the marked graph X , with respect to the sequence $\{(X_n, w_n)\}_{n \in \mathbb{N}}$, is defined by

$$\ln \zeta_{(X, w)}(z) = \lim_{n \rightarrow \infty} \frac{1}{|X_n|} \ln \zeta_{X_n}(z) = \sum_{r=1}^{\infty} \frac{\tilde{c}_r z^r}{r}, \quad |z| < \frac{1}{k-1}.$$

The proof that the series has a non-trivial radius of convergence is given in [8] and depends on results from [17].

Let X be a graph such that the degrees of vertices is bounded. Let Γ be a group of graph automorphisms of the graph X that acts on X without inversions and satisfying the following properties:

- (1) For each $v \in V(X)$, the stabilizer $\Gamma_v = \{\gamma \in \Gamma : \gamma v = v\}$ is finite.
- (2) If $\mathcal{F}_0 \subset V(X)$ is a complete set of orbit representatives of the action of Γ on $V(X)$, then

$$\text{vol}(X/\Gamma) = \sum_{v \in \mathcal{F}_0} \frac{1}{|\Gamma_v|} < \infty.$$

In particular, if the action of Γ on $V(X)$ is free, the second condition is equivalent to the condition that the orbit space $V(X)/\Gamma$ is finite. In this case, the Ihara zeta function is defined as

$$\zeta_{X, \Gamma}(z) = \prod_{C \in \mathcal{P}/\Gamma} \left(1 - z^{\ell(C)}\right)^{|\Gamma_C|}.$$

where:

- $\mathcal{P}$ are equivalence classes of closed, primitive, tail-less edge-paths without backtracking. Two such circuits are equivalent if they differ only by a shift. $\mathcal{P}/\Gamma$ denotes the orbit space of $\mathcal{P}$ under the Γ action.
- For each class $C \in \mathcal{P}/\Gamma$, $\ell(C)$ denotes the length of C i.e., the number of edges in C .
- Γ_C denotes the isotropy group of C .

This formula generalizes the classical zeta function on finite graphs.

We will describe the analogue of Bass' formula for $\zeta_{X, \Gamma}(z)$. Let $L^2(X)$ be the Hilbert space of functions on $V(X)$. A unitary representation is given by:

$$\lambda_0 : \Gamma \rightarrow U(L^2(X)), \quad (\lambda_0(\gamma)f)(v) = f(\gamma^{-1}v), \quad \gamma \in \Gamma, \quad f \in L^2(X), \quad v \in V(X).$$

Then the von Neumann algebra of all bounded operators on $L^2(X)$ that commute with the Γ action is defined as:

$$\mathcal{N}_0(X, \Gamma) = \{\lambda_0(\gamma) : \gamma \in \Gamma\}'.$$

The algebra $\mathcal{N}_0(X, \Gamma)$ inherits a trace given by:

$$\mathrm{Tr}_\Gamma(A) = \sum_{v \in \mathcal{F}_0} \frac{1}{|\Gamma_v|} A(v, v), \quad A \in \mathcal{N}_0(X, \Gamma).$$

With this setting, the Bass formula for the Ihara zeta function has the form ([3], [4], [10], [11], [12]):

$$\zeta_{X, \Gamma}^{-1}(z) = (1 - z^2)^{-\chi^{(2)}(X)} \det_\Gamma(\Delta_{X, z}),$$

where

- $\det_\Gamma = \exp \mathrm{Tr}_\Gamma \circ \ln$ is the determinant in the von Neumann algebra $\mathcal{N}_0(X, \Gamma)$.
- $\Delta_{X, z} = I - Az + Qz^2$, with A the adjacency operator on X , and Q is the operator on $L^2(X)$ given by:

$$Q(f)(v) = (\deg(v) - 1)f(v), \quad \text{for each } v \in V(X).$$

Remark 2.9.

- (1) In [3], [4], [10], [11], [12], it was shown that the function $\zeta_{X, \Gamma}$ is defined for sufficiently small $|u|$. More precisely, if k is the maximum degree of X , $\zeta_{X, \Gamma}(u)$ is a holomorphic function for all $|u| < \frac{1}{d-1}$.
- (2) $\chi^{(2)}(X)$ is the Euler characteristic defined in [3]. In most applications, it is equal to $\chi(X/\Gamma)$, the Euler characteristic of the orbit space.
- (3) Let X be a k -regular graph and $q = k - 1$. Using the determinant formula, the zeta function can be extended to a holomorphic function in the open set ([3], [10]):

$$\Omega_q = \mathbb{R}^2 \setminus \left(\left\{ (x, y) \in \mathbb{R}^2 : x^2 + y^2 = \frac{1}{q} \right\} \cup \left\{ (x, 0) \in \mathbb{R}^2 : \frac{1}{q} \leq |x| \leq 1 \right\} \right).$$

- (4) In the above references there is an interpretation of the Bass formula over the determinant on $\mathcal{N}_1(X, \Gamma)$, the von Neumann algebra on the set of edges of X .

Notation. There are three types of zeta functions used in this paper.

- (1) We write $\zeta_X(z)$ for the classical zeta function defined for a finite graph X .
- (2) We write $\zeta_{X, \Gamma}(z)$ for the equivariant zeta function defined on an infinite graph X equipped with an action of a group Γ with finite co-volume.
- (3) We write $\zeta_{(X, w)}(u)$ for the zeta function that it is the limit of $\zeta_{X_n}(z)^{1/|V(X_n)|}$, where $\{(X_n, w_n)\}_{n \in \mathbb{N}}$ is a covering sequence of finite regular graphs converging to (X, w) .

Definition 2.10. The sequence $\{X_n, w_n\}_{n \in \mathbb{N}}$ strongly converges to (X, w) if:

- (1) $\{X_n, w_n\}_{n \in \mathbb{N}}$ is a covering sequence of marked k -regular graphs with

$$p_{m-1} : X_m \rightarrow X_{m-1}$$

the covering map.

- (2) X is k -regular.
 (3) There are covering maps

$$\rho_n : X \rightarrow X_n$$

such that:

- (a) $\rho_1(u) = p_1 p_2 \dots p_{n-1}(\rho_n(u))$, for all n .
 (b) For each n , the isometry between $B_X(w, s_n)$ and $B_{X_n}(w_n, s_n)$ is given by the restriction of ρ_n .

Remark 2.11. Cayley graphs of groups give sequences of graphs that strongly converge. Let Γ be a group, S a symmetric set of generators and $\{K_n\}_{n \in \mathbb{N}}$ a sequence of normal subgroups of finite index such that:

$$K_1 \supset K_2 \supset K_3 \dots, \quad \text{and} \quad \bigcap_{n \in \mathbb{N}} K_n = \{K\}.$$

Then the sequence of the marked Schreier graphs $\{(\mathcal{S}(\Gamma, K_n, S), 1)\}_{n \in \mathbb{N}}$ strongly converges to $(\mathcal{S}(\Gamma, K, S), 1)$.

Let $\{X_n, w_n\}_{n \in \mathbb{N}}$ strongly converge to (X, w) and $\Gamma_n = \text{Cov}(X, X_n)$. The next results gives a connection between the different types of zeta functions.

Theorem 2.12. *Assume that all the graphs in the sequence are k -regular finite graphs converging to a k -regular graph (X, w) . Then*

$$\lim_{n \rightarrow \infty} \zeta_{X_n}(z)^{\frac{|V(X_1)|}{|V(X_n)|}} = \zeta_{X, \Gamma_1}(z) = \zeta_{(X, w)}(z)^{|V(X_1)|}.$$

Proof. By [4], Theorem 2.1,

$$\lim_{n \rightarrow \infty} \zeta_{X_n}(z)^{\frac{1}{|N_n|}} = \zeta_{X, \Gamma_1}(z)$$

where $N_n = [\Gamma_n, \Gamma_1] = |V(X_n)|/|V(X_1)|$. The result follows from the definition of $\zeta_{(X, w)}(z)$. $\square$

The following is the main part of the proof of Theorem 2.1 in [4].

Corollary 2.13. *With the above notation,*

$$\det_{\Gamma_1}(\Delta_{X, z}) = \lim_{n \rightarrow \infty} (\Delta_{X_n, z})^{\frac{|V(X_1)|}{|V(X_n)|}}$$


Let F be a locally finite graph and ϕ_1 an $\text{Aut}(F)$ -voltage assignment on X_1 . Inductively, define an $\text{Aut}(F)$ -voltage assignment on X_n by:

$$\phi_n(uv) = \phi_{n-1}(p_{n-1}(u)p_{n-1}(v)).$$

Also, define an $\text{Aut}(F)$ -voltage assignment on X by:

$$\phi(uv) = \phi_1(\rho_1(u)\rho_1(v)).$$

The details are presented in the following diagram.



Now, by Theorem 2.6, we know that, for any finite, d -regular graph F , the sequence $\{X_n \times^{\phi_n} F\}_{n \in \mathbb{N}}$ is a kd -regular covering sequence; thus, by [8], it converges. We will show that in fact it converges to the graph $X \times^{\phi} F$. To do this will need the following:

Lemma 2.14. *Assume that $\{(X_n, w_n)\}_{n \in \mathbb{N}}$ strongly converges to (X, w) . Then*

$$\widetilde{\rho}_n : B_{X \times^{\phi} F}((w, i), s_n) \rightarrow B_{X_n \times^{\phi_n} F}((w_n, i), s_n), \quad \widetilde{\rho}_n(u, i) = ((\rho_n(u), i),$$

is an isometry, for any $i \in V(F)$ and for all $n \in \mathbb{N}$.

Proof. Since ρ_n is a bijection, it is clear that $\widetilde{\rho}_n$ is a bijection; thus, we must show that $\widetilde{\rho}_n$ preserves adjacency. To this end, assume $(u, i) \sim (v, j)$, for $(u, i), (v, j) \in B_{X \times^\phi F}(w, i)$. Then there are two cases:

- (1) $u \sim v$ and $j = i^{\phi(uv)}$
- (2) $u = v$ and $i \sim j$.

In the case where $u = v$, since clearly $\rho_n(u) = \rho_n(v)$, we must have $\widetilde{\rho}_n(u, i) \sim \widetilde{\rho}_n(v, j)$. In the case where $u \sim v$, we must have $\rho_n(u) \sim \rho_n(v)$. Thus we must show that

$$j = i^{\phi_n(\rho_n(u)\rho_n(v))}.$$

Now, by the definition of ϕ ,

$$\phi(uv) = \phi_1(\rho_1(u)\rho_1(v)),$$

and by assumption,

$$\phi_1(\rho_1(u)\rho_1(v)) = \phi_1(p_{n-1} \dots p_1(\rho_n(u)), p_{n-1} \dots p_1(\rho_n(v))).$$

But by the definition of ϕ_n ,

$$\phi_n(\rho_n(u)\rho_n(v)) = \phi_1(p_{n-1} \dots p_1(\rho_n(u)), p_{n-1} \dots p_1(\rho_n(v))).$$

This shows that

$$\phi_n(\rho_n(u)\rho_n(v)) = \phi(uv),$$

and thus

$$j = i^{\phi(uv)} \implies j = i^{\phi_n(\rho_n(u)\rho_n(v))}.$$

This shows that $\hat{\rho}_n$ preserves adjacency, and thus is an isometry. This completes the proof. $\square$

As a corollary, we have the following theorem.

Theorem 2.15. *For each $i \in F$, the covering sequence $\{(X_n \times^{\phi_n} F, (w_n, i))\}_{n \in \mathbb{N}}$ strongly converges to $(X \times^\phi F, (w, i))$.*

Proof. Theorem 2.6 implies that the covering conditions of the strong convergence are satisfied. The rest of the proof follows from Lemma 2.14 and [8]. $\square$

3. ZETA FUNCTIONS FOR BUNDLES AND COVERINGS

In this section we will use our previous result and [3], [4], [10], [11], [12], to generalize the results of [6] to infinite graph bundles.

Definition 3.1. Let X be a graph equipped with an $\text{Aut}(F)$ -voltage assignment ϕ .

- (1) An action of a group Γ on X without edge inversions is called (F, ϕ) -compatible if

$$\phi(\gamma(u)\gamma(v)) = \phi(uv), \text{ for all } u, v \in V(X), \gamma \in \Gamma.$$

- (2) An action of a group Δ without inversions on F is called (X, ϕ) -compatible if $\text{Im}(\phi) \subset C_{\text{Aut}(F)}(\Delta)$ i.e., the image of ϕ centralizes Δ .
- (3) The pair of groups (Γ, Δ) as before is called ϕ -compatible if Γ is (F, ϕ) -compatible and Δ is (X, ϕ) -compatible.

Lemma 3.2. *With the above notation, if the pair (Γ, Δ) is ϕ -compatible, then the product action:*

$$(\gamma, \delta)(x, i) = (\gamma x, \delta i), \quad (\gamma, \delta) \in \Gamma \times \Delta, \quad (x, i) \in V(X \times^\phi F),$$

is an action by graph automorphisms on $X \times^\phi F$. Furthermore, if the action of Γ and Δ are of finite co-volume, so is the action of $\Gamma \times \Delta$ on $X \times^\phi F$.

Proof. The proof follows from the definitions. □

Theorem 3.3. *Assume that (Γ, Δ) is a pair of ϕ -compatible actions. Also, assume that the actions are of finite co-volume. Then*

$$\zeta_{X \times^\phi F, \Gamma \times \Delta}(z)^{-1} = (1 - z^2)^{-\chi^{(2)}(X \times^\phi F)} \det_{\Gamma \times \Delta} \left(I - \sum_{\gamma \in \text{Aut}(F)} (A_{\vec{X}_{(\phi, \gamma)}} \otimes P_\gamma + I_X \otimes A_F) z + Q z^2 \right),$$

$\chi^{(2)}$ is the Euler characteristic and Q is the diagonal operator with (x, i) -entry $\deg(x) + \deg(i) - 1$. Furthermore, the zeta function is holomorphic for $|z| < \frac{1}{k+d-1}$. If X is k regular and F d -regular, then $\zeta_{X \times^\phi F, \Gamma}(z)$ can be extended to a holomorphic function on Ω_{k+d-1} .

Proof. From [10], [11], we have

$$\zeta_{X \times^\phi F, \Gamma \times \Delta}(z)^{-1} = (1 - z^2)^{-\chi^{(2)}(X \times^\phi F)} \det_{\Gamma \times \Delta} (I - z A_{X \times^\phi F} + z^2 Q_{X \times^\phi F}).$$

The theorem now follows immediately from Theorem 2.7. □

We will now use Theorem 3.3 to provide a decomposition for the zeta function of any infinite cover. Let $p : Y \rightarrow X$ be a cover with X finite and Y locally finite. Let $\text{Cov}(p) = \Gamma$. Now we define the function

$$\phi : E(\vec{X}) \rightarrow \Gamma.$$

For this we write $X = \{x_1, \dots, x_n\}$. For each i , choose $v_i \in Y$ such that $p(v_i) = x_i$. Now, since $p : N(v_i) \rightarrow N(x_i)$ is a bijection, for each $x_j \in N(x_i)$ there exists a unique $u_j \in N(v_i)$ such that $p(u_j) = x_j$. So, since $p(v_j) = x_j = p(u_j)$, there exists some $\gamma \in \Gamma$ such that $\gamma v_j = u_j$. Thus, define

$$\phi : E(\vec{X}) \rightarrow \Gamma, \quad \phi(x_i x_j) = \gamma.$$

We then have the following:

Lemma 3.4. *Let ϕ be defined as above. Then*

- (1) *The Γ action on Y is of finite co-volume and it is ϕ -compatible.*

(2) The map

$$\alpha : Y \rightarrow X \times^\phi \Gamma, \quad \alpha(u) = (p(u), \beta)$$

is an isomorphism, where $\beta u = v_i$, $p(u) = x_i = p(v_i)$.

Proof. The proof is folklore. $\square$

Now, in order to prove an analogue of 2.7 for $Y \simeq X \times^\phi \Gamma_1$, we need to define the following operator: for $\gamma_1, \gamma_2, \gamma \in \Gamma_1$,

$$P_\gamma(\gamma_1, \gamma_2) = \begin{cases} 1, & \text{if } \gamma_2 = \gamma_1 \gamma \\ 0, & \text{otherwise.} \end{cases}$$

Lemma 3.5. *Let ϕ be defined as above. Then*

$$A_{X \times^\phi \Gamma} = \sum_{\gamma \in \Gamma} A_{\vec{X}_{(\phi, \gamma)}} \otimes P_\gamma.$$

Proof. The proof is analogous to that of 2.7. $\square$

The following theorem provides a decomposition for the zeta function of any infinite cover.

Theorem 3.6. *With the above notation,*

$$\zeta_{Y, \Gamma}(z)^{-1} = (1 - z^2)^{-\chi^{(2)}(X)} \det_\Gamma \left(I - \left(\sum_{\gamma \in \Gamma} A_{\vec{X}_{(\phi, \gamma)}} \otimes P_\gamma \right) z + Q z^2 \right),$$

where Q is the diagonal operator with (x, γ) -entry $\deg(x) - 1$. The function is holomorphic for $|z| < \frac{1}{k-1}$. If Y is k -regular then $\zeta_{Y, \Gamma}(z)$ is holomorphic on Ω_{k-1} .

Proof. This follows immediately from 3.3 and 3.4. $\square$

Let $\{(X_n, w_n)_{n \in \mathbb{N}}$ be a sequence of finite k -regular marked graphs that strongly converges to the k -regular marked graph (X, w) . Let F be a finite d -regular graph. With the notation as in Theorem 2.15, we know that

We write $a_n = |V(X_n)|$ and $f = |F|$.

Corollary 3.7. *With the above notation, for $|z| < \frac{1}{k+d-1}$,*

$$\begin{aligned} \zeta_{(X \times^\phi F, (w, i))}(z)^{-1} &= \\ &= \lim_{n \rightarrow \infty} \left[(1 - z^2)^{-\chi(X_n \times^\phi F)} \det \left(I - \sum_{\gamma \in \text{Aut}(F)} (A_{\vec{X}_n(\phi_n, \gamma)} \otimes P_\gamma + I_{X_n} \otimes A_F) z + Q_n z^2 \right) \right]^{\frac{1}{f a_n}} \\ &= \left[(1 - z^2)^{-\chi^{(2)}(X \times^\phi F)} \det_\Gamma \left(I - \sum_{\gamma \in \text{Aut}(F)} (A_{\vec{X}(\phi, \gamma)} \otimes P_\gamma + I_X \otimes A_F) z + Q z^2 \right) \right]^{\frac{1}{f a_1}} \end{aligned}$$

Proof. The first identity follows because of Theorem 2.15:

$$\begin{aligned} \zeta_{(X \times^\phi F, (w, i))}(z)^{-1} &= \lim_{n \rightarrow \infty} \zeta_{X_n \times^{\phi_n} F}(z)^{-\frac{1}{f a_n}} \\ &= \lim_{n \rightarrow \infty} \left[(1 - z^2)^{-\chi(X_n \times^{\phi_n} F)} \det \left(I - \sum_{\gamma \in \text{Aut}(F)} (A_{\overrightarrow{X_n}(\phi_n, \gamma)} \otimes P_\gamma + I_{X_n} \otimes A_F) z + Q_n z^2 \right) \right]^{\frac{1}{f a_n}} \end{aligned}$$

The second identity follows from Theorem 3.3 and Theorem 2.12. $\square$

4. APPLICATION

Let $p : Y \rightarrow X$ be a cover with $\mathbf{Cov}(p) = \Gamma$ and X a finite graph. let F be a finite d -regular graph with n such that $\text{Aut}(F)$ contains a the dihedral group D_{2n} of order $2n$. Let ϕ be an $\text{Aut}(F)$ -voltage assignment on X whose image is contained into D_{2n} and ψ the induced $\text{Aut}(F)$ -voltage assignment on Y (Theorem 2.6). By Theorem 2.6, the induced map

$$\tilde{p} : Y \times^\psi F \rightarrow X \times^\phi F, \quad \tilde{p}(x, i) = (p(x), i)$$

is a covering map. Also, $\text{Cov}(\tilde{p}) = \Gamma$.

The following is the setup (for the finite case this is the same as in [14] and [6]): set $V(F) = \{1, 2, \dots, n\}$ and S_n the symmetric group on $V(F)$. Let $a = (1 \ 2 \ \dots \ n-1 \ n)$ be an n -cycle and let

$$b = \begin{cases} (1 \ n)(2 \ n-1) \dots \left(\frac{n-1}{2} \ \frac{n+3}{2} \right) \left(\frac{n+1}{2} \right) & \text{if } n \text{ is odd,} \\ (1 \ n)(2 \ n-1) \dots \left(\frac{n}{2} \ \frac{n+2}{2} \right) & \text{if } n \text{ is even} \end{cases}$$

be a permutation in S_n . The permutations a and b generate the dihedral subgroup D_n of S_n :

$$D_n = \langle a, b \mid a^n = b^2 = 1, bab = a^{-1} \rangle.$$

Let $\mu = \exp(2\pi i/n)$ and $\mathbf{x}_k = (1, \mu^k, \mu^{2k}, \dots, \mu^{(n-1)k})^T$ be the column vector in $\mathbb{C}^n$. Then $1, \mu, \dots, \mu^{n-1}$ are the distinct eigenvalues of the permutation matrix $P(a)$ and $\mathbf{x}_k$ is the eigenvector corresponding to the eigenvalue μ^k . Let $P(b)$ be the permutation matrix of b and

$$M = \begin{cases} [\mathbf{x}_0 & P(b)\mathbf{x}_1 & x_2 & P(b)\mathbf{x}_2 & \dots & x_{(n-1)/2} & P(b)x_{(n-1)/2}] & \text{if } n \text{ is odd} \\ [\mathbf{x}_0 & P(b)\mathbf{x}_1 & x_2 & P(b)\mathbf{x}_2 & \dots & x_{(n-2)/2} & P(b)x_{(n-2)/2} & \mathbf{x}_{n/2}] & \text{if } n \text{ is even} \end{cases}$$

In [14], (also [6]) it was shown that $P(b)\mathbf{x}_k$ is an eigenvector of $P(a)$ associated with the eigenvalue μ^{n-k} . Thus M is invertible. Also, $P(a)$ and A_F commute and thus they are simultaneously diagonalizable. Also, $\mathbf{x}_k$ and $P(b)\mathbf{x}_k$ ($1 \leq k \leq (n-1)/2$ when n is odd and $1 \leq k \leq (n-2)/2$ when n is even) are eigenvectors of A_F with the same eigenvalue of $P(b)$, denoted $\lambda_{(F,k)}$. Also, $\mathbf{x}_0$ is

the eigenvector of A_F corresponding to the eigenvalue d and, for n even, $\lambda_{(F,n/2)}$ is the eigenvalue associated to the eigenvector $\mathbf{x}_2$. Then as in [14], using Theorem 2.7, we get that

$$(I_Y \otimes M)^{-1} A_{Y \times \psi F} (I_Y \otimes M) = \begin{cases} (A_Y + dI_Y) \oplus \left(\bigoplus_{i=1}^{(n-1)/2} (A_t + \lambda_{(F,t)} (I_Y \oplus I_Y)) \right) & \text{if } n \text{ is odd} \\ (A_Y + dI_Y) \oplus \left(\bigoplus_{i=1}^{(n-2)/2} (A_t + \lambda_{(F,t)} (I_Y \oplus I_Y)) \oplus (B + \lambda_{(F,n/2)} I_Y) \right) & \text{if } n \text{ is even} \end{cases}$$

where

$$B = \sum_{k=0}^{n-1} \left((-1)^k A(\vec{Y}_{(\psi, a^k)}) + (-1)^{k+1} A(\vec{Y}_{(\psi, a^k b)}) \right),$$

and

$$A_t = \sum_{k=0}^{n-1} \begin{bmatrix} \mu^{tk} A(\vec{Y}_{(\psi, a^k)}) & \mu^{tk} A(\vec{Y}_{(\psi, a^k b)}) \\ \mu^{(n-t)k} A(\vec{Y}_{(\psi, a^k b)}) & \mu^{(n-t)k} A(\vec{Y}_{(\psi, a^k)}) \end{bmatrix}$$

Also, let $L_Y = (Q_Y + dI_Y) \otimes I_2$. Then the calculation in section 4 in [6] can be carried through in our setting and we get the following:

Theorem 4.1. *Let $p : Y \rightarrow X$ be as above. Then*

$$\zeta_{Y \times \psi F, \Gamma}(z)^{-1} = (1 - z^2)^{-\chi^{(2)}(Y \times \psi F)} f_{Y, F}(z) \prod_{t=1}^{(n-1)/2} g_{Y, F, t}(z)$$

when n is odd, and

$$\zeta_{Y \times \psi F, \Gamma}(z)^{-1} = (1 - z^2)^{-\chi^{(2)}(Y \times \psi F)} f_{X, F}(z) h_{Y, F}(z) \prod_{t=1}^{(n-2)/2} g_{Y, F, t}(z)$$

when n is even, where

- (1) $g_{Y, F, t}(z) = \det_{\Gamma} (I_Y \oplus I_Y - (A_t + \lambda_{(F, t)} (I_Y \oplus I_Y)) z + L_Y z^2)$
- (2) $h_{Y, F}(z) = \det_{\Gamma} (I_Y - (B + \lambda_{(F, n/2)} I_Y) z + (Q_Y + dI_Y) z^2)$
- (3) $f_{X, F}(z) = \det_{\Gamma} (I_Y - (A_Y + dI_Y) z + (Q_Y + dI_Y) z^2)$

Proof. This follows by simple calculation from 3.3 with $\Delta = \{1\}$, and Theorem 9 of [6]. $\square$

Let $\{(X_m, w_m)\}_{m \in \mathbb{N}}$ be a sequence of finite regular graphs that strongly converges to (X, w) . Let F be a finite d -regular with n vertices such that $\text{Aut}(F)$ contains D_{2n} . Let ϕ be an $\text{Aut}(F)$ -voltage assignment on X_1 whose image is contained in D_{2n} . Let ϕ_n be the induced $\text{Aut}(F)$ -voltage on X_m and ψ be the induced $\text{Aut}(F)$ -voltage assignment on X . Set

$$\Gamma = \text{Cov}(X \rightarrow X_1), \quad \Delta_m = \text{Cov}(X_m \rightarrow X_1), \quad m \in \mathbb{N}.$$

Corollary 4.2. *Let $a_m = |V(X_m)|$. With the above notation,*

(1) *If n is odd:*

$$\zeta_{X \times^\psi F, \Gamma}(z)^{-1} = \zeta_{(X \times^\psi F, (w, i))}(z)^{-a_1} = \lim_{m \rightarrow \infty} \left((1 - z^2)^{-\chi(X_m \times^{\phi_m} F)} f_{X_m, F}(z) \prod_{t=1}^{(n-1)/2} g_{X_m, F, t}(z) \right)^{\frac{a_1}{a_m}}.$$

(2) *If n is even*

$$\zeta_{X \times^\psi F, \Gamma}(z)^{-1} = \zeta_{(X \times^\psi F, (w, i))}(z)^{-a_1} = \lim_{m \rightarrow \infty} \left((1 - z^2)^{-\chi(X_m \times^{\phi_m} F)} f_{X_m, F}(z) h_{X_m, F}(z) \prod_{t=1}^{(n-1)/2} g_{X_m, F, t}(z) \right)^{\frac{a_1}{a_m}}.$$

Proof. It follows from Theorem 2.12, Theorem 2.15 and Theorem 4.1. $\square$

In some cases, we can get a better description of the functions appearing in the expression for the zeta function of the limit. Assume that all the graphs X_m , $m \in \mathbb{N}$, and X are p -regular. Following [8], for each $m \in \mathbb{N}$ set:

$$\sigma_m = \sum \frac{\delta_{\lambda_i(X_m)}}{a_m}$$

where $\lambda_i(X_m)$ are the eigenvalues of the Markov operator $(1/k)A_{X_m}$ on X_m and δ_x is the Dirac function. The sequence $\{\sigma_m\}_{m \in \mathbb{N}}$ weakly converges to the spectral measure σ associated to the Markov operator $(1/k)A_X$. Using the calculation of Section 5 in [8] and Corollary 2.13, we get:

$$\begin{aligned} \ln f_{X, F}(z) &= \ln \det_{\Gamma} (I_Y - (A_Y + dI_Y)z + (Q_Y + dI_Y)z^2) \\ &= \lim_{m \rightarrow \infty} \frac{1}{a_m} \ln \det (I_{X_m} - (A_{X_m} + dI_{X_m})z + (Q_{X_m} + dI_{X_m})z^2) \\ &= \lim_{m \rightarrow \infty} \int_{-1}^1 \ln (1 - (p\lambda + d)z + (p - 1 + d)z^2) d\sigma_m(\lambda) \\ &= \int_{-1}^1 \ln (1 - (p\lambda + d)z + (p - 1 + d)z^2) d\sigma(\lambda) \end{aligned}$$

where σ is the spectral measure associated to $(1/p)A_X$.

Summarizing:

Corollary 4.3. *With the above notation,*

$$\ln f_{X, F}(z) = \int_{-1}^1 \ln (1 - (p\lambda + d)z + (p - 1 + d)z^2) d\sigma(\lambda), \text{ for } |z| < \frac{1}{p + d - 1},$$

where σ is the spectral measure associated to the regular random walk on X .

We give a specific example. The same method works for any group for which the spectral measure is known. Let Γ be the Grigorchuk group ([1], [5], [7], [8]). Then Γ can be represented as a subgroup of automorphisms of the rooted binary tree. Let $P = \text{St}(1^\infty)$ be the stabilizer of the

infinite sequence of 1's. Let P_m be the stabilizer of all the elements that start with m 1's and it has finite index in Γ . Then

$$P = \bigcup_{m=1}^{\infty} P_m.$$

If $S = \{a, b, c, d\}$ be the standard set of generators of Γ , then the Schreier graphs $\{\mathbb{S}_m = \mathcal{S}(\Gamma, P_m, S)\}_{m \in \mathbb{N}}$ converge to $\mathbb{S} = \mathcal{S}(\Gamma, P, S)$. All the graphs have as the base point the identity coset. Then in [8], Corollary 9.2, we have that

$$\begin{aligned} \ln \zeta_{\mathbb{S}, P}(z) &= -3 \ln(1 - z^2) - \int_{-1/2}^0 \frac{(1 - 8xz + 7z^2) |1 - 4x|}{2\pi \sqrt{x(2x-1)(2x+1)(1-x)}} dx \\ &\quad - \int_{1/2}^1 \frac{(1 - 8xz + 7z^2) |1 - 4x|}{2\pi \sqrt{x(2x-1)(2x+1)(1-x)}} dx \end{aligned}$$

Let F be a d -regular graph as in the beginning of the section and ϕ an $\text{Aut}(F)$ -voltage assignment whose image lies into D_{2n} . Combining Theorem 4.1 and Corollary 4.3 we get.

Theorem 4.4. *With the above notation, let ψ be the $\text{Aut}(F)$ -voltage assignment on $\mathbb{S}$ and $R = P_1/P$. Then*

$$\begin{aligned} &\ln \zeta_{(\mathbb{S} \times^{\psi} F, (P, i))}(z) \\ &= \frac{1}{da_1} \left[-\chi^{(2)}(\mathbb{S} \times^{\phi} F) \ln(1 - z^2) + \ln \det_R \left(I - \sum_{\gamma \in \text{Aut}(F)} (A(\vec{\mathbb{S}}_{(\psi, \gamma)}) \otimes P_{\gamma} + I_{\mathbb{S}} \otimes A_F) z + Q z^2 \right) \right] \end{aligned}$$

Furthermore,

$$\zeta_{(\mathbb{S} \times^{\psi} F, (P, i))}(z)^{-1} = (1 - z^2)^{-\chi^{(2)}(\mathbb{S} \times^{\psi} F)} f_{\mathbb{S}, F}(z) \prod_{t=1}^{(n-1)/2} g_{\mathbb{S}, F, t}(z)$$

when n is odd, and

$$\zeta_{(\mathbb{S} \times^{\psi} F, (P, i))}(z)^{-1} = (1 - z^2)^{-\chi^{(2)}(\mathbb{S} \times^{\psi} F)} f_{X, F}(z) h_{\mathbb{S}, F}(z) \prod_{t=1}^{(n-2)/2} g_{\mathbb{S}, F, t}(z)$$

when n is even, where g and h are as in Theorem 4.1 and

$$\ln f_{\mathbb{S}, F}(z) = \int_{-\frac{1}{2}}^0 \frac{\ln(1 - (8x + d)z + (7 + d)z^2) |1 - 4x|}{2\pi \sqrt{x(2x-1)(2x+1)(1-x)}} dx + \int_{\frac{1}{2}}^1 \frac{\ln(1 - (8x + d)z + (7 + d)z^2) |1 - 4x|}{2\pi \sqrt{x(2x-1)(2x+1)(1-x)}} dx$$

REFERENCES

- [1] L. Bartholdi, R. I. Grigorchuk, *On the spectrum of Hecke type operators related to some fractal groups*, Proc. Steklov Inst. Math., **231** (2000), 1–41.
- [2] H. Bass, *The Ihara–Selberg zeta function of a tree lattice*, Internat. J. Math. **3** (1992), 717–797.
- [3] B. Clair, S. Mokhtari-Sharghi, *Zeta functions of discrete groups acting on trees*, J. Algebra **237** (2001), 591–430.
- [4] B. Clair, S. Mokhtari-Sharghi, *Convergence of zeta functions of graphs*, Proc. Amer. Math. Soc. **130** (2002), 1881–1886.

- [5] P. de la Harpe, *Topics in geometric group theory*, Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 2000.
- [6] R. Feng, J. Kwak, K. Kim, *Zeta functions of graph bundles*, preprint.
- [7] R. I. Grigorchuk, V. V. Nekrashevich, V. I. Sushchanskii, *Automata, dynamical systems, and groups*, Proc. Steklov Inst. Math., **231** (2000), 128–203.
- [8] R. Grigorchuk, A. Żuk, *The Ihara zeta function of infinite graphs, the KNS spectral measure and integrable maps*, in: “Random Walks and Geometry”, Proc. Workshop (Vienna 2001), V. A. Kaimanovich *et al.*, eds., de Gruyter, Berlin 2004, 141–180.
- [9] R. Grigorchuk, A. Żuk, *On the asymptotic spectrum of random walks on infinite families of graphs*, Random walks and discrete potential theory (Cortona, 1997), 188–204, Sympos. Math., XXXIX, Cambridge Univ. Press, Cambridge, 1999.
- [10] D. Guido, T. Isola, M. L. Lapidus, *Ihara zeta functions for periodic simple graphs*, arXiv:math.OA/0605753, May 2006.
- [11] D. Guido, T. Isola, M. L. Lapidus, *Ihara zeta functions for periodic simple graphs and its approximation in the amenable case*, arXiv:math.OA/0608229, August 2006.
- [12] D. Guido, T. Isola, M. L. Lapidus, *A trace on fractal graphs and the Ihara zeta function*, arXiv:math.OA/0608060, August 2006.
- [13] Y. Ihara, *On discrete subgroups of the two by two projective linear group over p -adic fields*, J. Math. Soc. Japan **18** (1966), 219–235.
- [14] J. H. Kwak, Y. S. Kwon, *Characteristic polynomials of graph bundles having voltages in a dihedral group*, Linear Algebra Appl. **336** (2001), 99–118.
- [15] J. H. Kwak, J. Lee, *Characteristic polynomials of some graph bundles II*, Linear and Multilinear Algebra **32** (1992), 61–73.
- [16] A. Lubotzky, *Discrete Groups, Expander Graphs and Invariant Measures*
- [17] J.-P. Serre, *Répartitions asymptotique des valeurs propres de l’opérateur de Hecke T_p* , J. Amer. Math. Soc. **10** (1997), 75–102.
- [18] H. M. Stark, A. A. Terras, *Zeta functions of finite graphs and coverings*, Adv. Math. **121** (1996), 126–165.
- [19] H. M. Stark, A. A. Terras, *Zeta functions of finite graphs and coverings II*, Adv. Math. **154** (2000), 132–195.
- [20] H. M. Stark, A. A. Terras, *Zeta functions of finite graphs and coverings III*, Adv. Math.

DEPARTMENT OF MATHEMATICS, VANDERBILT UNIVERSITY, NASHVILLE, TN 37240

E-mail address: samuel.d.cooper@vanderbilt.edu

DEPARTMENT OF MATHEMATICS CANISIUS COLLEGE, BUFFALO, NY 14208

E-mail address: prasside@canisius.edu